Recombinant Human B7-DC/PD-L2/CD273 Protein

Catalog No.: RP00150

www.abclonal.com

Sequence Information

Species
Human

Gene ID
80380

Swiss Prot
Q9BQ51

Tags
C-His

Synonyms
B7DC,Btdc,CD273,PD-L2, PDCD1L2,PDL2,bA574F11.2, PDCD1LG2

Product Information

Source
HEK293 cells

Purification
≥ 95 % as determined by SDS-PAGE;≥ 90 % as determined by HPLC.

Calculated MW
23.40 kDa

Observed MW
35-50 kDa

Endotoxin
< 0.1 EU/μg of the protein by LAL method.

Formulation
Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Programmed Death Ligand 2 (PD-L2), also known as B7-DC and butyrophilin-like protein, is a member of the B7 family of proteins that provide signals for regulating T-cell activation and tolerance .PD-L2 is expressed on dendritic cells, subsets of activated CD4+ and CD8+ T cells, and memory B cells that differentiate into plasma cells. At inflammatory sites such as rheumatoid arthritis, allergen exposure, and virus infection, PD-L2 is up-regulated on synoviocytes, infiltrating macrophages, dendritic cells, and airway epithelial cells. PD-L2, along with B7-H1/PD-L1, binds to T cell PD-1 where it promotes IFN-gamma production and CD40 Ligand up-regulation while inhibiting IL-4 production. In addition, PD-L2 binds to RGM-B on macrophages and alveolar epithelial cells, supporting respiratory immune tolerance. In asthma, PD-L2 suppresses IL-5 and IL-13 production, promotes IL-12 production by dendritic cells, and supports allergen-induced airway hyper-responsiveness and mucus production.

Basic Information

Description
Recombinant Human B7-DC/PD-L2/CD273 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Leu20-Pro219) of human B7-DC/PD-L2/CD273 (Accession #NP_079515.2) fused with a 6×His tag at the C-terminus.

Activity
Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human PD-L2 at 5 μg/mL (100 μL/well) can bind Recombinant Human PD-1 with a linear range of 0.3-1.2 μg/mL.

Shipping
Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage
Store at -20℃


Operational Notes
For your safety and health, please wear a lab coat and disposable gloves for handling.

Contact


400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.


Validation Data



Immobilized Recombinant Human PD-L2 at 5μg/mL (100 μL/well) can bind Recombinant Human PD-1 with a linear range of 0.3-1.2 μg/mL.



The purity of Human B7-DC/PD-L2/CD273 Protein (Cat.RP00150) was greater than 90% as determined by SEC-HPLC.



Recombinant Human B7-DC/PD-L2/CD273 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.