

Recombinant Human Fc-gamma RIII beta/FCGR3B /CD16b Protein

Catalog No.: RP00471

Sequence Information

Species	Gene ID
Human	2215
Swiss Prot	Tags
O75015-1	C-His&Avi

Synonyms

CD16; Fc gamma RIIB; FCG3; FCGR3B; FcgRIIB; FCRIIB; IGFR3; CD16b (NA2); CD16B; FCG3B; FCG3; FCGR3; FCR-10

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE; ≥ 95 % as determined by HPLC.

Calculated MW Observed MW

23.7 kDa 47-53 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Human Fc gamma RIIB/CD16b Protein is a receptor for the Fc region of immunoglobulins gamma. Low affinity receptor. Binds complexed or aggregated IgG and also monomeric IgG. Contrary to III-A, is not capable to mediate antibody-dependent cytotoxicity and phagocytosis. May serve as a trap for immune complexes in the peripheral circulation which does not activate neutrophils.

Basic Information

Description

Recombinant Human Fc-gamma RIII beta/FCGR3B/CD16b Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly17-Ser200) of Human Fc-gamma RIII beta/FCGR3B /CD16b (Accession #O75015-1) fused with a C-His&Avi tag at the C-terminus.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature. Liquid proteins are shipped at frozen temperature with dry ice. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Recombinant Human Fc-gamma RIII beta/FCGR3B/CD16b Protein was determined by Tris-Bis PAGE under reducing conditions.



Human Fc gamma RIIIB (NA2), His Tag captured on CM5 Chip via Anti-his antibody can bind Rituximab with an affinity constant of 2.33 μ M as determined in SPR assay (Biacore T200).



The purity of Human Fc gamma RIIIB (NA2) is greater than 95% as determined by SEC-HPLC.