

Recombinant Human Fc gamma RIIA/FCGR2A /CD32a (H167R) Protein

Catalog No.: RP00480

Sequence Information

Species	Gene ID
Human	2212
Swiss Prot	Tags
P12318-1	C-His&Avi

Synonyms

Fc gamma RIIA; FCG2; FCGR2;
FCGR2A; FCGR2A1; FcγRIIA;
FCRIIA; fcRII-a; FCG2; CD32A

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE; ≥ 95 % as determined by HPLC.

Calculated MW	Observed MW
23.2 kDa	34-40 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

The Fc gamma Rs have been divided into three classes based on close relationships in their extracellular domains; these groups are designated Fc gamma RI (also known as CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type. The CD64 proteins are high affinity receptors (10^{-8} - 10^{-9} M) capable of binding monomeric IgG, whereas the CD16 and CD32 proteins bind IgG with lower affinities (10^{-6} - 10^{-7} M) only recognizing IgG aggregates surrounding multivalent antigens.

Basic Information

Description

Recombinant Human Fc gamma RIIA/FCGR2A/CD32a (H167R) Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala36-Ile218(H167R)) of Human Fc gamma RIIA/CD32a (Accession #P12318-1) fused with a C-His&Avi tag at the C-terminus.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature. Liquid proteins are shipped at frozen temperature with dry ice. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

		400-999-6126
		cn.market@abclonal.com.cn
		www.abclonal.com.cn


Validation Data



Recombinant Human Fc gamma RIIA/FCGR2A/CD32a (H167R) Protein was determined by Tris-Bis PAGE under reducing conditions.



Human Fc gamma RIIA (R167) , His Tag captured on CM5 Chip via anti-his antibody can bind Rituximab with an affinity constant of 2.89 μ M as determined in SPR assay (Biacore T200).



The purity of Human Fc gamma RIIA (R167) is greater than 95% as determined by SEC-HPLC.