

Biotinylated Recombinant Human CTLA-4/CD152 Protein

Catalog No.: RP00505B

Sequence Information

Species	Gene ID
Human	1493
Swiss Prot	Tags
P16410	C-His&Avi

Synonyms

CTLA4; CD152; ICOS; CELIAC3;
GRD4; GSE; ALPS5; IDDM12

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE; ≥ 95 % as determined by HPLC.

Calculated MW	Observed MW
16.3 kDa	25-30 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

CTLA-4 (cytotoxic T-lymphocyte-associated protein 4), also known as CD152, is a protein receptor that, functioning as an immune checkpoint, downregulates immune responses. CTLA4 is constitutively expressed in regulatory T cells but only upregulated in conventional T cells after activation – a phenomenon which is particularly notable in cancers. It acts as an "off" switch when bound to CD80 or CD86 on the surface of antigen-presenting cells.

Basic Information

Description

Biotinylated Recombinant Human CTLA-4/CD152 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Lys36-Asp161) of Human CTLA-4/CD152 (Accession #P16410) fused with a C-His&Avi tag at the C-terminus.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature. Liquid proteins are shipped at frozen temperature with dry ice. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

		400-999-6126
		cn.market@abclonal.com.cn
		www.abclonal.com.cn

Validation Data




Biotinylated Recombinant Human CTLA-4/CD152 Protein was determined by Tris-Bis PAGE under reducing conditions.




The purity of Biotinylated Human CTLA-4 is greater than 95% as determined by SEC-HPLC.

Biotinylated Human CTLA-4, His Tag ELISA
0.1 µg Biotinylated Human CTLA-4, His Tag Per Well



Immobilized Biotinylated Human CTLA-4, His Tag at 1 µg/mL (100 µL/well) on the streptavidin precoated plate (5 µg/mL). Dose response curve for Human B7-1, hFc Tag with the EC_{50} of 2.4ng/mL determined by ELISA (QC Test).



Immobilized Anti-CTLA-4 Antibody, hFc Tag at 1 µg/mL (100 µL/well) on the plate. Dose response curve for Biotinylated Human CTLA-4, His Tag with the EC_{50} of 45.7 ng/mL determined by ELISA.