Biotinylated Recombinant Human TNFRSF17 /BCMA/CD269 Protein

Catalog No.: RP00612B

www.abclonal.com

Sequence Information

Species
Human

Gene ID
608

Swiss Prot
Q02223-1

Tags
C-His&Avi

Synonyms
CD269; TNFRSF17; BCMA; BCM; TNFRSF13A

Product Information

Source
HEK293 cells

Purification
≥ 95 % as determined by SDS-PAGE;≥ 95 % as determined by HPLC.

Calculated MW
8.9 kDa

Observed MW
12, 15-17 kDa

Endotoxin
< 1 EU/μg of the protein by LAL method.

Formulation
Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

B-cell maturation antigen (BCMA or BCM), also known as tumor necrosis factor receptor superfamily member 17 (TNFRSF17), is a protein that in humans is encoded by the TNFRSF17 gene.TNFRSF17 is a cell surface receptor of the TNF receptor superfamily which recognizes B-cell activating factor (BAFF).

Basic Information

Description
Biotinylated Recombinant Human TNFRSF17/BCMA/CD269 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Ala54) of Human TNFRSF17/BCMA/CD269 (Accession #Q02223-1) fused with a C-His&Avi tag at the C-terminus.

Activity

Shipping
Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage
Store at -20℃


Operational Notes
For your safety and health, please wear a lab coat and disposable gloves for handling.

Contact


400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.


Validation Data



Immobilized Anti-BCMA Antibody, hFc Tag at 0.2 μg/mL (100 μL/Well) on the plate. Dose response curve for Biotinylated Human BCMA, His Tag with the EC_{50} of 3.4ng /mL determined by ELISA.



The purity of Biotinylated Human BCMA is greater than 95% as determined by SEC-HPLC.



Biotinylated Recombinant Human TNFRSF17/BCMA/CD269 Protein was determined by Tris-Bis PAGE under reducing conditions.