

Recombinant Human CEACAM6/CD66c Protein

Catalog No.: RP00982

Sequence Information

Species Gene ID

Human 4680

Swiss Prot Tags

P40199 C-His

Synonyms

CEACAM6, NCA, Cell adhesion molecule CEACAM6, Carcinoembryonic antigen-related cell adhesion molecule 6, CEA cell adhesion molecule 6, Non-specific crossreacting antigen, Normal cross-reacting antigen, CD66c

Product Information

Source Purification

HEK293 cells ≥ 95 % as determined by SDS-PAGE.

Calculated MW Observed MW

32.09 kDa 40-65 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Basic Information

Description

Recombinant Human CEACAM6/CD66c Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Lys35-Gly320) of human CEACAM6 (Accession #NP_002474.3) fused with a 6×His tag at the C-terminus.

Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human CEACAM6 at 2 μg/mL (100 μL/well) can bind Recombinant Human CEACAM8 with a linear range of 0.1-0.4 μg/mL.

Shipping

Lyophilized proteins are shipped at ambient temperature. Liquid proteins are shipped at frozen temperature with dry ice. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Immobilized Recombinant Human CEACAM6 at 2 $\mu\text{g/mL}$ (100 μL /well) can bind Recombinant Human CEACAM8 with a linear range of 0.1-0.4 $\mu\text{g/mL}$.



Recombinant Human CEACAM6/CD66c Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.