

Recombinant Human Transferrin/TF Protein

Catalog No.: RP00997

Sequence Information

Species Gene ID

Human 7018

Swiss Prot Tags

P02787 C-His

Synonyms

HEL-S-71p,PRO1557,PRO2086,
TFQTL1,TF,Transferrin

Product Information

Source Purification

HEK293 cells
≥ 95 % as
determined by
SDS-PAGE; ≥95
% as determined
by HPLC.

Calculated MW Observed MW

76.02 kDa 76-90 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL
method.

Formulation

Lyophilized from a 0.22 μm filtered
solution of PBS, pH 7.4. Contact us
for customized product form or
formulation.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled
water. Avoid vortex or vigorously
pipetting the protein. For long
term storage, it is recommended
to add a carrier protein or stabilizer
(e.g. 0.1% BSA, 5% HSA, 10% FBS
or 5% Trehalose), and aliquot the
reconstituted protein solution to
minimize free-thaw cycles.

Background

Basic Information

Description

Recombinant Human Transferrin/TF Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val20-Pro698) of human Transferrin (Accession #NP_001054.1) fused with a 6×His tag at the C-terminus.

Activity

1. Measured in a serum-free cell proliferation assay using MCF-7 human breast cancer cells. The ED₅₀ for this effect is 0.075-0.3 ng/mL.
2. Measured by its binding ability in a functional ELISA. Immobilized Recombinant human Transferrin at 2 μg/mL (100 μL/well) can bind Recombinant human TfR with a linear range of 4-17 ng/mL.

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data




Recombinant Human Transferrin/TF Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



The purity of Human Transferrin Protein (Cat.RP00997) was greater than 95% as determined by SEC-HPLC.



Immobilized Recombinant human Transferrin at 2 $\mu\text{g/mL}$ (100 $\mu\text{L/well}$) can bind Recombinant human TfR with a linear range of 4–17 ng/mL.



Recombinant Human Transferrin promotes the proliferation of MCF-7 human breast cancer cells. The ED_{50} for this effect is 0.075–0.3 ng/mL.