

Recombinant Human Cystatin-D/CST5 Protein

Catalog No.: RP01035

Sequence Information

Species	Gene ID
Human	1473
Swiss Prot	Tags
P28325	C-His

Synonyms

CST5, MGC71922, CST5

Product Information

Source	Purification
HEK293 cells	≥ 90 % as determined by SDS-PAGE.

Calculated MW Observed MW

14.70 kDa 15-25 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Cystatins are natural inhibitors of papain-like (family C1) and legumain-related (family C13) cysteine peptidases. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. As a member of type 2 cystatin, cystatin D is a single-domain protein and also has cysteine residues that form disulfide bridges. The functions of Cystatin D are largely unknown. However, Cystatin D has been shown to inhibit coronavirus replication at its physiological concentration (0.12_x005f

Basic Information

Description

Recombinant Human Cystatin-D/CST5 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly21-Val142) of human Cystatin D (Accession #NP_001891.2) fused with a 6×His tag at the C-terminus.

Activity

Measured by its ability to inhibit papain cleavage of a fluorogenic peptide substrate Z-FR-AM. The IC50 value is <18 nM.

Shipping

Lyophilized proteins are shipped at ambient temperature. Liquid proteins are shipped at frozen temperature with dry ice. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Recombinant Human Cystatin-D/CST5 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Human CST6 inhibits papain cleavage of a fluorogenic peptide substrate Z-FR-AMC (Catalog # ES009). The IC₅₀ value is <18 nM.