

Recombinant Human ACE-2 Protein

Catalog No.: RP01266

Sequence Information

Species	Gene ID
Human	59272
Swiss Prot	Tags
Q9BYF1	C-His&Avi

Synonyms

ACE2, ACEH, angiotensin-converting enzyme 2, ACEH

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE; ≥95 % as determined by HPLC.

Calculated MW Observed MW

86.25 kDa 95-110 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. or Supplied as a 0.22 μm filtered solution in PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Basic Information

Description

Active Recombinant Human ACE2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln18-Ser740) of human ACE2 (Accession #Q9BYF1) fused with a 6×His, Avi tag at the C-terminus.

Activity

Measured by its binding ability in a functional ELISA. Immobilized Human ACE2 at 2 μg/mL (100 μL/well) can bind SARS-CoV-2 Spike RBD with a linear range of 0.03-4.9 ng/mL.

Shipping

Lyophilized proteins are shipped at ambient temperature. Liquid proteins are shipped at frozen temperature with dry ice. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data




Recombinant Human ACE-2 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



SARS-COV-2 Spike S1, hFc Tag captured on CM5 Chip via Protein A can bind Human ACE2, His Tag with an affinity constant of 3.80 nM as determined in SPR assay (Biacore T200).



Immobilized Human ACE2 at 2 $\mu\text{g/mL}$ (100 μL /well) can bind SARS-CoV-2 Spike RBD with a linear range of 0.03–4.9 ng/mL.



The purity of Human ACE2 is greater than 95% as determined by SEC-HPLC