

Recombinant Human CD200R1 Protein

Catalog No.: RP01365

Sequence Information

Species Gene ID

Human 131450

Swiss Prot Tags
Q8TD46 C-hFc&His

Synonyms

CD200R1, CD200R, HCRTR2, MOX2R, OX2R

Product Information

Source Purification

HEK293 cells ≥ 95 % as determined by SDS-PAGE.

Calculated MWObserved MW

53.59 kDa 70-100 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cell surface glycoprotein CD200 receptor 1 (CD200R1) is an isoform of CD200 receptors that is expressed on cells of the myeloid lineage. CD200R1 is a receptor for the OX-2 membrane glycoprotein. The receptor-substrate interaction may serve as a myeloid downregulatory signal.

Basic Information

Description

Recombinant Human CD200R1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala27-Leu266) of human CD200R/MOX2R (Accession #NP_620161.1) fused with a Fc, 6×His tag at the C-terminus.

Activity

Measured by its binding ability in a functional ELISA. Immobilized Human CD200 at 2 μg/mL (100 μL/well) can bind Human CD200 R with a linear range of 0.058-8.34 ng/mL.

Shipping

Lyophilized proteins are shipped at ambient temperature. Liquid proteins are shipped at frozen temperature with dry ice. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Immobilized Human CD200 at 2 $\mu\text{g/mL}$ (100 $\mu\text{L/well}$) can bind Human CD200 R with a linear range of 0.058-8.34 ng/mL.



Recombinant Human CD200R1 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.