Recombinant Mouse ICOS/CD278 Protein

Catalog No.: RP01572

Sequence Information

Species
Mouse

Gene ID
54167

Swiss Prot
Q9WVS0

Tags
C-hFc

Synonyms
ICOS,CD278,AILIM,CVID1,ICOS

Product Information

Source
HEK293 cells

Purification
≥ 95 % as determined by SDS-PAGE.

Calculated MW
40.15 kDa

Observed MW
55-60 kDa

Endotoxin
< 0.1 EU/μg of the protein by LAL method.

Formulation
Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Inducible costimulator (ICOS), also called AILIM (Activation-Inducible Lymphocyte Immunomediatory Molecule) is a cell-surface receptor and belongs to the CD28 family of immune costimulatory receptors consisting of CD28, CTLA-4, and PD-1. The interaction of B7-H2/ICOS plays a critical role in Th cell differentiation, T/B cell interactions which are essential for the germinal center formation, and humoral immune responses, and as well as the production of cytokine IL-4. Also, ICOS is more potent in the induction of IL-10 production, a cytokine important for the suppressive function of T regulatory cells. The B7-1/B7-2--CD28/CTLA-4 and ICOS-B7RP-1 pathway provide key second signals that can regulate the activation, inhibition, and fine-tuning of T-lymphocyte responses. ICOS stimulates both Th1 and Th2 cytokine production but may have a preferential role in Th2 cell development. Moreover, The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway has been suggested as being involved in the development of airway inflammation and airway hyperresponsiveness.

Basic Information

Description
Recombinant Mouse ICOS/CD278 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu21-Leu144) of mouse ICOS/CD278 (Accession #NP_059508.2) fused with a hFc tag at the C-terminus.

Activity

Shipping
Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage
Store at -20℃


Operational Notes
For your safety and health, please wear a lab coat and disposable gloves for handling.

Contact


400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.

Validation Data



Immobilized Mouse ICOSLG at 5 μg/mL (100 μL/well) can bind Mouse ICOS with a linear range of 10-952.7 ng/mL



Recombinant Mouse ICOS/CD278 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.