

Recombinant Mouse CCL2/MCP-1 Protein

Catalog No.: RP01607

Sequence Information

Species Gene ID

Mouse 20296

Swiss Prot Tags

P10148 C-His

Synonyms

C-C motif chemokine ligand 2,
CCL2,GDCF-2,HC11,HSMCR30,
MCAF,Mcp1,MCP-1,SCYA2,SMC-
CF,CCL2

Product Information

Source Purification

HEK293 cells ≥ 95 % as
determined by
SDS-PAGE; ≥ 95
% as determined
by HPLC.

Calculated MW Observed MW

14.69 kDa 25-37 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL
method.

Formulation

Lyophilized from a 0.22 μm filtered
solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled
water. Avoid vortex or vigorously
pipetting the protein. For long
term storage, it is recommended
to add a carrier protein or stabilizer
(e.g. 0.1% BSA, 5% HSA, 10% FBS
or 5% Trehalose), and aliquot the
reconstituted protein solution to
minimize free-thaw cycles.

Background

Basic Information

Description

Recombinant Mouse CCL2/MCP-1 Protein is produced by HEK293 cells
expression system. The target protein is expressed with sequence (Gln 24-
Asn148) of mouse CCL2/MCP-1 (Accession #NP_035463.1) fused with a
6×His tag at the C-terminus.

Activity

Measured by its ability to chemoattract THP-1 cells. The ED₅₀ for this effect is
typically 6.2-25.0 ng/mL.

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for
handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Recombinant Mouse CCL2/MCP-1 Protein is greater than 95% as determined by SEC-HPLC.



Recombinant Mouse CCL2/MCP-1 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Mouse CCL2 chemoattract THP-1 cells. The ED₅₀ for this effect is typically 6.2-25.0 ng/mL.