

Recombinant human Trk-A/NTRK1 Protein

Catalog No.: RP01726

Sequence Information

Species **Gene ID**
human 4914

Swiss Prot **Tags**
P04629-1 C-6His

Synonyms

MTC; TRK; TRK1; TRKA; Trk-A;
p140-TrkA;NTRK1

Product Information

Source **Purification**
HEK293 cells ≥ 90 % as
 determined by
 SDS-PAGE.

Calculated MW **Observed MW**
42.35 kDa 60-100 kDa

Endotoxin

< 0.01 EU/μg of the protein by LAL
method

Formulation

Lyophilized from a 0.22 μm filtered
solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled
water. Avoid vortex or vigorously
pipetting the protein. For long
term storage, it is recommended
to add a carrier protein or stabilizer
(e.g. 0.1% BSA, 5% HSA, 10% FBS
or 5% Trehalose), and aliquot the
reconstituted protein solution to
minimize free-thaw cycles.

Background

This protein is a member of the neurotrophic tyrosine kinase receptor (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. The presence of this kinase leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in this gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. Alternate transcriptional splice variants of this gene have been found, but only three have been characterized to date.

Basic Information

Description

Recombinant human Trk-A/NTRK1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala33-Glu413) of human Trk-A/NTRK1 (Accession #NP_001012331.1) fused with and a 6×His tag at the C-terminus.

Activity

Measured by its ability to inhibit NGF-induced proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is 0.086-0.34 μg/mL in the presence of 10 ng/mL of Recombinant Human beta-NGF.

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Recombinant human Trk-A/NTRK1 inhibit NGF-induced proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is 0.086-0.34 μ g/mL in the presence of 10 ng/mL of Recombinant Human beta-NGF.



Recombinant human Trk-A/NTRK1 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.