

# Recombinant Human DLL1 Protein

**Catalog No.: RP02151**

## Sequence Information

**Species**      **Gene ID**  
Human            28514

**Swiss Prot**    **Tags**  
O00548          C-His

**Synonyms**  
DLL1, DELTA1, DL1, Delta

## Product Information

**Source**            **Purification**  
HEK293 cells    ≥ 95 % as  
determined by  
SDS-PAGE.

**Calculated MW** **Observed MW**  
57.3 kDa          65 kDa

**Endotoxin**  
< 1 EU/μg of the protein by LAL  
method.

## Formulation

Lyophilized from a 0.2 μm filtered  
solution of 20mM Hepes, 150mM  
NaCl, 1mM EDTA, pH 7.4. Contact  
us for customized product form or  
formulation.

## Reconstitution

Centrifuge the tube before  
opening. Reconstitute to a  
concentration of 0.1-0.5 mg/mL in  
sterile distilled water. Avoid vortex  
or vigorously pipetting the protein.  
For long term storage, it is  
recommended to add a carrier  
protein or stabilizer (e.g. 0.1% BSA,  
5% HSA, 10% FBS or 5%  
Trehalose), and aliquot the  
reconstituted protein solution to  
minimize free-thaw cycles.

## Background

Delta-like protein 1 (DLL1) is a type I transmembrane protein that belongs to the Delta/Serrate/Lag2 (DSL) family of Notch ligands. Mature human DLL1 consists of a 528 amino acid (aa) extracellular domain (ECD) with one DSL domain and eight EGF-like repeats, a 23 aa transmembrane segment, and a 155 aa cytoplasmic domain. Within the ECD, human DLL1 shares 91% aa sequence identity with mouse and rat DLL1. The residual membranebound portion of DLL1 can be cleaved by presenilin-dependent γ-secretase, enabling the cytoplasmic domain to migrate to the nucleus. DLL1 localizes to adherens junctions on neuronal processes through its association with the scaffolding protein MAGI1. DLL1 is widely expressed, and it plays an important role in embryonic somite formation, cochlear hair cell differentiation, plus B and T lymphocyte differentiation. The upregulation of DLL1 in arterial endothelial cells following injury or angiogenic stimulation is central to postnatal arteriogenesis. DLL1 is also overexpressed in cervical carcinoma and glioma and contributes to tumor progression.

## Basic Information

### Description

Recombinant Human DLL1 Protein is produced by Mammalian expression system. The target protein is expressed with sequence (Gln18-Gly540) of human DLL1 (Accession #O00548) fused with a 6xHis tag at the C-terminus.

### Activity

### Shipping

Lyophilized proteins are shipped at ambient temperature.  
Liquid proteins are shipped at frozen temperature with dry ice.  
Upon receipt, store it immediately at the temperature recommended below.

### Storage

Store at -20°C

### Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

## Contact


☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

**Validation Data**

---



Recombinant Human DLL1 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.