

Biotinylated Recombinant Human ROR2 Protein

Catalog No.: RP02480B

Sequence Information

Species	Gene ID
Human	4920
Swiss Prot	Tags
A1L4F5(Q01974)	C-His&Avi

Synonyms

BDB, BDB1, NTRKR2, ROR2, BDB1, NTRKR2

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by Tris-Bis PAGE; ≥ 95 % as determined by HPLC.

Calculated MW Observed MW

44.2 kDa 54-65 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

ROR2 (Receptor Tyrosine Kinase-like Orphan Receptor 2) is a member of the ROR family of receptor tyrosine kinases and is important for skeletal development, including bone and cartilage formation, as well as for the development of the central nervous system. Mature human ROR2 contains a 369 amino acid (aa) extracellular domain (ECD) and a 518 aa cytoplasmic tail containing an tyrosine kinase domain.

Basic Information

Description

Biotinylated Recombinant Human ROR2 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Val34-Gly403) of Human ROR2 fused with a His and Avi tag at the C-terminal.

Activity

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Immobilized Anti-ROR2 Antibody, hFc Tag at 0.5µg/ml (100µl/well) on the plate.
Dose response curve for Biotinylated Human ROR2, hFc Tag with the EC₅₀ of 6.5ng /ml determined by ELISA.



The purity of Biotinylated Human ROR2 is greater than 95% as determined by SEC-HPLC.



Biotinylated Recombinant Human ROR2 Protein was determined by Tris-Bis PAGE under reducing conditions.