Recombinant Human PRCP Protein

Catalog No.: RP03204LQ

Sequence Information

Species
Human

Gene ID
5547

Swiss Prot
P42785

Tags
C-His

Synonyms
Lysosomal Pro-X Carboxypeptidase; Angiotensinase C; Lysosomal Carboxypeptidase C; Proline Carboxypeptidase; Prolylcarboxypeptidase; PRCP; PRCP; PCP

Product Information

Source
HEK293 cells

Purification
≥ 95 % as determined by SDS-PAGE.

Calculated MW
54.5 KDa

Observed MW
60-90 KDa

Endotoxin
< 1 EU/μg of the protein by LAL method.

Formulation
Supplied as a 0.2 μm filtered solution of 20mM NaAc-HAc, 150mM NaCl, 10% Glycerol, pH4.5.

Reconstitution

Background

Lysosomal Pro-X Carboxypeptidase (PRCP) belongs to the peptidase S28 family. PRCP is detected in many tissues, with highest levels observed in placenta, lung, and liver. It is also present in the heart, brain, pancreas, and kidney. PRCP exists as a homodimer. PRCP cleaves C-terminal amino acids linked to proline in peptides such as angiotensin II, III and des-Arg9-bradykinin. This cleavage occurs at acidic pH, but enzymatic activity is retained with some substrates at neutral pH. PRCP has been shown to be an activator of the cell matrix-associated prekallikrein.

Basic Information

Description
Recombinant Human PRCP Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu22-His496) of Human PRCP (Accession #NP_005031.1) fused with 6×His tag at the C-terminus.

Activity

Shipping
Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage
Store at -70℃


Operational Notes
For your safety and health, please wear a lab coat and disposable gloves for handling.

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.

Validation Data



Recombinant Human PRCP Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.