Recombinant Human EGF Protein

Catalog No.: RP03287

www.abclonal.com

Sequence Information

Species
Human

Gene ID
1950

Swiss Prot
P01133

Tags
No-Tag

Synonyms
EGF,Pro-epidermal growth factor, EGF, Cleaved into: Epidermal growth factor, Urogastrone

Product Information

Source
E. coli

Purification
≥ 95 % as determined by SDS-PAGE.≥ 95 % as determined by HPLC.

Calculated MW
6.35 kDa

Observed MW
5-15 kDa

Endotoxin
< 0.01 EU/μg of the protein by LAL method

Formulation
Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

As a low-molecular-weight polypeptide, EGF was first purified from the mouse submandibular gland, but since then it was found in many human tissues including submandibular gland, parotid gland. It can also be found in human platelets, macrophages, urine, saliva, milk, and plasma. EGF is a growth factor that stimulates the growth of various epidermal and epithelial tissues in vivo and in vitro and of some fibroblasts in cell culture. It results in cellular proliferation, differentiation, and survival. Salivary EGF, which seems also regulated by dietary inorganic iodine, also plays an important physiological role in the maintenance of oro-esophageal and gastric tissue integrity. EGF acts by binding with high affinity to epidermal growth factor receptor on the cell surface and stimulating the intrinsic protein-tyrosine kinase activity of the receptor.

Basic Information

Description
Recombinant Human EGF Protein is produced by E. coli Cells expression system. The target protein is expressed with sequence (Asn971-Arg1023) of Human EGF(Accession #NP_001954.2) fused with no tag .

Activity
Measured in a cell proliferation assay using Balb/C 3T3 mouse embryonic fibroblasts. The ED50 for this effect is typically 0.02-0.2ng/ml.

Shipping
Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage
Store at -20℃


Operational Notes
For your safety and health, please wear a lab coat and disposable gloves for handling.

Contact


400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.


Validation Data




Recombinant Human EGF Protein is greater than 95% as determined by SEC-HPLC.




Human kidney organoids were cultured with EGF(Cat. RP03287), FGF2(Cat. RP01042), FGF7(Cat. RP01717), FGF9(Cat. RP01710), FGF10(Cat. RP01140), IGF-(Cat. RP00996), NOG(Cat. RP01237), RSPO1(Cat. RP00071), WNT-3a(Cat. RP01618SLQ). And further, DKK-1(RP01343) was used to induce the establishment of cell polarity.




Recombinant Human EGF Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.




Mouse lung organoids were cultured in Matrigel (Cat. RPM0003) with medium supplemented with EGF (Cat. RP03287), Noggin (Cat. RP01237), RSPO1 (Cat. RP00071), Wnt-3a (Cat. RP01618SLQ), and FGF-10 (Cat. RP01140).




Human stomach organoids were cultured with EGF(Cat. RP03287), FGF10(Cat. RP01140), NOG(Cat. RP01237), RSPO1(Cat. RP00071), WNT-3a(Cat. RP01618SLQ).




Measured in a cell proliferation assay using Balb3T3 mouse fibroblast cells. The ED_{50} for this effect is 0.04-0.17 ng/mL, corresponding to a specific activity of 5.88×10^{6}~2.50×10^{7} units/mg.



Human liver organoids were cultured with EGF(Cat. RP03287), HGF(Cat. RP01602), FGF2(Cat. RP01042), FGF10(Cat. RP01140), NOG(Cat. RP01237), RSPO1(Cat. RP00071), WNT-3a(Cat. RP01618SLQ).



Human colorectal cancer organoid were cultured with EGF(Cat. RP03287), NOG(Cat. RP01237)， FGF10(Cat. RP01140), RSPO1(Cat. RP00071), HGF(Cat. RP01602).



Human gastric cancer organoid were cultured with EGF(Cat. RP03287), NOG(Cat. RP01237), FGF10(Cat. RP01140), RSPO1(Cat. RP00071), Wnt-3a Surrogate(Cat. RP01618SLQ).