

Recombinant Human Proto-oncogene tyrosine-protein kinase Src/SRC Kinase

Catalog No.: RP03356LQ

Sequence Information

Species **Gene ID**
Human 6714

Swiss Prot **Tags**
P12931 N-His-Flag

Synonyms

SRC; SRC1; p60-Src; Proto-oncogene c-Src; Proto-oncogene tyrosine-protein kinase Src

Product Information

Source **Purification**
Baculovirus-Insect Cells ≥ 90 % as determined by SDS-PAGE; ≥ 90 % as determined by HPLC.

Calculated MW **Observed MW**
61.6 kDa 50-60 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Supplied as a 0.22 μm filtered solution in 50 mM HEPES, 150 mM NaCl, 1 mM ATP, 5 mM MgCl₂ (pH 7.5). Contact us for customized product form or formulation.

Reconstitution

Please use running water to thaw it quickly.

Background

Proto-oncogene tyrosine-protein kinase Src, also known as c-Src, is a non-receptor tyrosine kinase protein that in humans is encoded by the SRC gene. It includes an SH2 domain, an SH3 domain and a tyrosine kinase domain. When src is activated, it promotes survival, angiogenesis, proliferation and invasion pathways. It also regulates angiogenic factors and vascular permeability after focal cerebral ischemia-reperfusion, and regulates matrix metalloproteinase-9 activity after intracerebral hemorrhage.

Basic Information

Description

Recombinant Human Proto-oncogene tyrosine-protein kinase Src/SRC Protein is produced by Baculovirus-Insect Cells expression system. The target protein is expressed with sequence (Gly2-Leu536) of Human SRC (Accession #P12931) fused with a N-His-Flag tag.

Activity

The activity of SRC is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -70°C

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Recombinant Human Proto-oncogene tyrosine-protein kinase Src/SRC Kinase was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.



The activity of SRC is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.