RXRα Rabbit mAb

Catalog No.: A19105

Recombinant | 10 Publications

Basic Information

Observed MW
51 kDa

Calculated MW
51 kDa

Category
Monoclonal Antibodies

Applications
WB, IP, IF/ICC, IHC-P, ChIP, ChIP-seq, ELISA

Cross Reactivity
Human, Mouse, Rat

Clone No.
ARC0468

Background

Retinoid X receptors (RXRs) and retinoic acid receptors (RARs) are nuclear receptors that mediate the biological effects of retinoids by their involvement in retinoic acid-mediated gene activation. These receptors function as transcription factors by binding as homodimers or heterodimers to specific sequences in the promoters of target genes. The protein encoded by this gene is a member of the steroid and thyroid hormone receptor superfamily of transcriptional regulators. Alternative splicing of this gene results in multiple transcript variants.

Recommended Dilutions

WB	1:2000 - 1:10000
IP	0.5 μg-4 μg antibody for 400 μg-600 μg extracts of whole cells
IF/ICC	1:100 - 1:1000
IHC-P	1:100 - 1:400
ChIP	5 μg antibody for 10 μg-15 μg of Chromatin
ChIP-seq	5 μg antibody for 25 μg of Chromatin
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
6256

Swiss-Prot
P19793

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
NR2B1; RXRalpha; RXR-alpha; RXRα

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG

Purity
Affinity purification

Storage
Store at -20℃


Validation Data




Chromatin immunoprecipitation was performed with 25 µg of cross-linked chromatin from Hep G2 cells using 5 µg of RXRα Rabbit mAb (A19105). DNA libraries were prepared using Scale ssDNA-seq Lib Prep Kit for Illumina V2 (RK20228). The ChIP sequencing results indicate the enrichment pattern of RXRα across chromosome 19 (upper panel) and the genomic region encompassing ECH1, a representative gene enriched in RXRα (lower panel).




Chromatin immunoprecipitation was performed with 25 µg of cross-linked chromatin from Hep G2 cells using 5 µg of RXRα Rabbit mAb (A19105). DNA libraries were prepared using Scale ssDNA-seq Lib Prep Kit for Illumina V2 (RK20228). The ChIP sequencing results indicate the enrichment pattern of RXRα in the representative genomic region surrounding ECH1 gene.




Western blot analysis of lysates from MCF7 cells using RXRα Rabbit mAb (A19105) at 1:5000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30 s.




Western blot analysis of various lysates using RXRα Rabbit mAb (A19105) at 1:5000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 45 s.




Chromatin immunoprecipitation analysis of extracts from Hep G2 cells, using RXRα Rabbit mAb (A19105) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Confocal imaging of HeLa cells using RXRα Rabbit mAb (A19105,dilution 1:100) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012,dilution 1: 400) (Green). DAPI was used for nuclear staining (blue). Objective: 100x.



Immunoprecipitation of RXRα from 600 µg extracts of Hep G2 cells was performed using 3 µg of RXRα Rabbit mAb (A19105). Rabbit IgG isotype control (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X reducing Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using RXRα Rabbit mAb (A19105) at a dilution of 1:500.



Immunohistochemistry analysis of paraffin-embedded Human colon tissue using RXRα Rabbit mAb (A19105) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.