Leader in Biomolecular Solutions for Life Science

www.abclonal.com

DDX4 Rabbit mAb

Catalog No.: A27291

Recombinant | 1 Publications

Basic Information

Observed MW
80 kDa

Calculated MW
79 kDa

Category
Monoclonal Antibodies

Applications
WB, IP, IF-F, IF-P, IHC-P, ELISA

Cross Reactivity
Human, Mouse, Rat

Clone No.
ARC72110

Background

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein, which is a homolog of VASA proteins in Drosophila and several other species. The gene is specifically expressed in the germ cell lineage in both sexes and functions in germ cell development. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB	1:5000 - 1:30000
IP	0.5μg-4μg antibody for 200μg-400μg extracts of whole cells
IF-F	1:200 - 1:800
IF-P	1:200 - 1:800
IHC-P	1:700 - 1:7000
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
54514

Swiss-Prot
Q9NQI0

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
VASA

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃


Validation Data




Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using DDX4 Rabbit mAb (A27291) at a dilution of 1:7000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat testis tissue using DDX4 Rabbit mAb (A27291) at a dilution of 1:7000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Confocal imaging of paraffin-embedded Mouse testis tissue using DDX4 Rabbit mAb (A27291, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of paraffin-embedded Rat testis tissue using DDX4 Rabbit mAb (A27291, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Western blot analysis of various lysates using DDX4 Rabbit mAb (A27291) at 1: 14000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Negative control (NC): Mouse brain
Exposure time: 10s.




Immunoprecipitation of DDX4 from 300 µg extracts of Mouse testis was performed using 0.5 µg of DDX4 Rabbit mAb (A27291). Rabbit IgG isotype control (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X reducing Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using DDX4 Rabbit mAb (A27291) at a dilution of 1:10000.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.



Immunohistochemistry analysis of paraffin-embedded Human testis tissue using DDX4 Rabbit mAb (A27291) at a dilution of 1:7000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of frozen sections of Mouse testis tissue using DDX4 Rabbit mAb (A27291, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.