

CD28 Rabbit mAb**Catalog No.: A30090****Recombinant****Basic Information****Observed MW**

14 kDa, 40-60 kDa

Calculated MW

6-26 kDa

Category

Monoclonal Antibodies

Applications

WB, IP, IF/ICC, IF-F, IF-P, IHC-P, ELISA

Cross Reactivity

Human, Mouse

Clone No.

ARC4107

Background

The protein encoded by this gene is essential for T-cell proliferation and survival, cytokine production, and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions**WB** 1:10000 - 1:25000**IP**

0.4 µg - 2 µg antibody for 200 µg - 400 µg extracts of whole cells

IF/ICC 1:400 - 1:1600**IF-F** 1:400 - 1:1600**IF-P** 1:400 - 1:1600**IHC-P** 1:1000 - 1:4000**ELISA**

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions ($\geq 1:10000$), a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information**Gene ID**

940

Swiss-Prot

P10747

Immunogen

This information is considered to be commercially sensitive.

Synonyms

Tp44; IMD123


Contact

☎		400-999-6126
✉		cn.market@abclonal.com.cn
🌐		www.abclonal.com.cn

Product Information

Source	Isotype	Purity
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C		

Validation Data



Western blot analysis of various lysates using CD28 Rabbit mAb (A30090) at 1:25000 dilution incubated overnight at 4°C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Negative control (NC): THP-1.


Exposure time: 45 s.




Immunoprecipitation of CD28 from 300 µg extracts of Jurkat clone A3 [A3] cells was performed using 0.4 µg of CD28 Rabbit mAb (A30090). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using CD28 Rabbit mAb (A30090) at a dilution of 1:50000.




Confocal imaging of frozen sections of Mouse spleen tissue using CD28 Rabbit mAb (A30090, dilution 1:1200) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Human tonsil tissue using CD28 Rabbit mAb (A30090, dilution 1:1200) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of RPMI 8226 cells and THP-1 cells (negative) using CD28 Rabbit mAb (A30090, dilution 1:1200) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of EL4 cells using CD28 Rabbit mAb (A30090, dilution 1:1200) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using CD28 Rabbit mAb (A30090) at a dilution of 1:2500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using CD28 Rabbit mAb (A30090) at a dilution of 1:2500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse thymus tissue using CD28 Rabbit mAb (A30090) at a dilution of 1:2500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.