Leader in Biomolecular Solutions for Life Science

www.abclonal.com

K48-linkage Specific Ubiquitin Rabbit mAb

Catalog No.: A3606

Recombinant | 20 Publications

Basic Information

Observed MW
17-250 kDa

Calculated MW
26 kDa

Category
Monoclonal Antibodies

Applications
WB, IF/ICC, DB, ELISA

Cross Reactivity
Human

Clone No.
ARC0811

Background

This gene encodes ubiquitin, one of the most conserved proteins known. Ubiquitin has a major role in targeting cellular proteins for degradation by the 26S proteosome. It is also involved in the maintenance of chromatin structure, the regulation of gene expression, and the stress response. Ubiquitin is synthesized as a precursor protein consisting of either polyubiquitin chains or a single ubiquitin moiety fused to an unrelated protein. This gene consists of three direct repeats of the ubiquitin coding sequence with no spacer sequence. Consequently, the protein is expressed as a polyubiquitin precursor with a final amino acid after the last repeat. An aberrant form of this protein has been detected in patients with Alzheimer's disease and Down syndrome. Pseudogenes of this gene are located on chromosomes 1, 2, 13, and 17. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB	1:1000 - 1:6000
IF/ICC	1:50 - 1:200
DB	1:1000 - 1:3000

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
7314

Swiss-Prot
P0CG47

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
HEL-S-50; K48-linkage Specific Ubiquitin

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃


Validation Data




Confocal imaging of MCF7 cells using K48-linkage Specific Ubiquitin Rabbit mAb (A3606,dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007,dilution 1:500)(Red).The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green).DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of HeLa cells (treated with MG-132) and HeLa cells (untreated) cells using K48-linkage Specific Ubiquitin Rabbit mAb (A3606, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Western blot analysis of lysates from HeLa cells using K48-linkage Specific Ubiquitin Rabbit mAb (A3606) at 1:1000 dilution incubated overnight at 4℃. HeLa cells were treated with MG132 (50 μM) at 37℃ for 90 minutes.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25μg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.



Dot-blot analysis of all sorts of peptides using K48-linkage Specific Ubiquitin Rabbit mAb (A3606) at 1:1000 dilution.