

Musashi-1 (MSI1) Rabbit mAb**Catalog No.: A9122****Recombinant****Basic Information****Observed MW**

39 kDa

Calculated MW

39 kDa

Category

Monoclonal Antibodies

Applications

WB, IP, IF/ICC, ELISA

Cross Reactivity

Human, Mouse

Clone No.

ARC1796

Background

This gene encodes a protein containing two conserved tandem RNA recognition motifs. Similar proteins in other species function as RNA-binding proteins and play central roles in posttranscriptional gene regulation. Expression of this gene has been correlated with the grade of the malignancy and proliferative activity in gliomas and melanomas. A pseudogene for this gene is located on chromosome 11q13.

Recommended Dilutions**WB** 1:500 – 1:2000**IP**

0.5 µg – 4 µg antibody for 200 µg – 400 µg extracts of whole cells

IF/ICC 1:50 – 1:200**ELISA**

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information**Gene ID**

4440

Swiss-Prot

O43347

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

MSI1; musashi RNA binding protein 1; Musashi-1 (MSI1)

Contact | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn**Product Information****Source**

Rabbit

Isotype

IgG


Purity

Affinity purification


Storage

Store at -20°C


Validation Data



Immunofluorescence analysis of NIH-3T3 cells using Musashi-1 (MSI1) Rabbit mAb (A9122) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using Musashi-1 (MSI1) Rabbit mAb (A9122) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using Musashi-1 (MSI1) Rabbit mAb (A9122) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3min.



Immunoprecipitation of Musashi-1 (MSI1) from 300 µg extracts of SH-SY5Y cells was performed using 2 µg of Musashi-1 (MSI1) Rabbit mAb (A9122). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Musashi-1 (MSI1) Rabbit mAb (A9122) at a dilution of 1:1000.