Leader in Biomolecular Solutions for Life Science

www.abclonal.com

[KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb

Catalog No.: A19068

Recombinant | 4 Publications

Basic Information

Observed MW
150 kDa

Calculated MW
129 kDa

Category
Monoclonal Antibodies

Applications
WB, IF-F, IF-P, IHC-P, ELISA

Cross Reactivity
Human, Mouse, Rat

Clone No.
ARC0457

Background

This gene encodes the alpha subunit of a transmembrane receptor for collagens and related proteins. The encoded protein forms a heterodimer with a beta subunit and mediates the adhesion of platelets and other cell types to the extracellular matrix. Loss of the encoded protein is associated with bleeding disorder platelet-type 9. Antibodies against this protein are found in several immune disorders, including neonatal alloimmune thrombocytopenia. This gene is located adjacent to a related alpha subunit gene. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB	1:1000 - 1:6000
IF-F	1:200 - 1:400
IF-P	1:200 - 1:400
IHC-P	1:200 - 1:800

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
3673

Swiss-Prot
P17301

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
BR; GPIa; CD49B; HPA-5; VLA-2; VLAA2; b)

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃


Validation Data




Immunohistochemistry analysis of paraffin-embedded Human breast tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Western blot analysis of lysates from wild type(WT) and Integrin alpha 2 (ITGA2 /CD49b) knockout (KO) 293T cells, using [KO Validated] Integrin alpha 2 (ITGA2 /CD49b) Rabbit mAb (A19068) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.




Western blot analysis of lysates from A549 cells, using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.




Western blot analysis of various lysates, using [KO Validated] Integrin alpha 2 (ITGA2 /CD49b) Rabbit mAb (A19068) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.




Confocal imaging of paraffin-embedded Human colon tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Human kidney tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Mouse colon tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Rat colon tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Rat kidney tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of frozen sections of Mouse colon tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of frozen sections of Rat colon tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of frozen sections of Rat kidney tissue using [KO Validated] Integrin alpha 2 (ITGA2/CD49b) Rabbit mAb (A19068, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.