Leader in Biomolecular Solutions for Life Science

USP11 Rabbit pAb

Catalog No.: A20179

www.abclonal.com

Basic Information

Observed MW
110kDa

Calculated MW
110kDa

Category
Polyclonal Antibodies

Applications
WB, IF/ICC, ELISA

Cross Reactivity
Human, Mouse, Rat

Background

Protein ubiquitination controls many intracellular processes, including cell cycle progression, transcriptional activation, and signal transduction. This dynamic process, involving ubiquitin conjugating enzymes and deubiquitinating enzymes, adds and removes ubiquitin. Deubiquitinating enzymes are cysteine proteases that specifically cleave ubiquitin from ubiquitin-conjugated protein substrates. This gene encodes a deubiquitinating enzyme which lies in a gene cluster on chromosome Xp11.23

Recommended Dilutions

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:200

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
8237

Swiss-Prot
P51784

Immunogen
This information is considered to be commercially sensitive.

Synonyms
UHX1; USP11

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃


Validation Data



Western blot analysis of various lysates using USP11 Rabbit pAb (A20179) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.



Western blot analysis of lysates from Rat testis, using USP11 Rabbit pAb (A20179) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Immunofluorescence analysis of U2OS cells using USP11 Rabbit pAb (A20179) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.