Leader in Biomolecular Solutions for Life Science

DiMethyl-Histone H3-K27 Rabbit pAb

Catalog No.: A2362

4 Publications

www.abclonal.com

Basic Information

Observed MW
17 kDa

Calculated MW
15 kDa

Category
Polyclonal Antibodies

Applications
WB, IP, IF/ICC, IHC-P, DB, ChIP, ChIP-seq, ELISA

Cross Reactivity
Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB	1:500 - 1:5000
IP	0.5 μg - 4 μg antibody for 400 μg - 600 μg extracts of whole cells|
IF/ICC	1:50 - 1:200
IHC-P	1:50 - 1:200
DB	1:500 - 1:1000
ChIP	5μg antibody for 5μg-10μg of Chromatin
ChIP-seq	1:20 - 1:100
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
8290/8350

Swiss-Prot
Q16695/P68431

Immunogen
This information is considered to be commercially sensitive.

Synonyms
H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; DiMethyl-Histone H3-K27

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃


Validation Data




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using DiMethyl-Histone H3-K27 Rabbit pAb (A2362) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using DiMethyl-Histone H3-K27 Rabbit pAb (A2362) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat lung tissue using DiMethyl-Histone H3-K27 Rabbit pAb (A2362) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Chromatin immunoprecipitation analysis of extracts of 293 cell line, using DiMethyl-Histone H3-K27 Rabbit pAb (A2362) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Dot-blot analysis of all sorts of methylation peptides using DiMethyl-Histone H3-K27 Rabbit pAb (A2362) at 1:1000 dilution.



Immunofluorescence analysis of HeLa cells using DiMethyl-Histone H3-K27 Rabbit pAb (A2362) at dilution of 1:20 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates, using DiMethyl-Histone H3-K27 Rabbit pAb (A2362) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.