CBFB Rabbit mAb

Catalog No.: A28470

Recombinant

Basic Information

Observed MW
22 kDa

Calculated MW
22 kDa

Category
Monoclonal Antibodies

Applications
WB, IF/ICC, IHC-P, FC (intra), ELISA

Cross Reactivity
Human, Mouse, Rat

Clone No.
ARC69665

Background

The protein encoded by this gene is the beta subunit of a heterodimeric core-binding transcription factor belonging to the PEBP2/CBF transcription factor family which master-regulates a host of genes specific to hematopoiesis (e.g., RUNX1) and osteogenesis (e.g., RUNX2). The beta subunit is a non-DNA binding regulatory subunit; it allosterically enhances DNA binding by alpha subunit as the complex binds to the core site of various enhancers and promoters, including murine leukemia virus, polyomavirus enhancer, T-cell receptor enhancers and GM-CSF promoters. Alternative splicing generates two mRNA variants, each encoding a distinct carboxyl terminus. In some cases, a pericentric inversion of chromosome 16 [inv(16)(p13q22)] produces a chimeric transcript consisting of the N terminus of core-binding factor beta in a fusion with the C-terminal portion of the smooth muscle myosin heavy chain 11. This chromosomal rearrangement is associated with acute myeloid leukemia of the M4Eo subtype. Two transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB	1:2000-1:5000
IF/ICC	1:50-1:200
IHC-P	1:200-1:800
FC (intra)	1:100-1:500

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.For high-ratio antibody dilutions (≥1:10000)，a sequential dilution method is strongly recommended to ensure measurement accuracy.For

Immunogen Information

Gene ID
865

Swiss-Prot
Q13951

Immunogen
Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms
CLCD2; PEBP2B

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃


Validation Data




Western blot analysis of various lysates using CBFB Rabbit mAb (A28470) at 1:5000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90 s.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using CBFB Rabbit mAb (A28470) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human spleen tissue using CBFB Rabbit mAb (A28470) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using CBFB Rabbit mAb (A28470) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse thymus tissue using CBFB Rabbit mAb (A28470) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat spleen tissue using CBFB Rabbit mAb (A28470) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Western blot analysis of lysates from K-562 cells using CBFB Rabbit mAb (A28470) at 1:5000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 45 s.



Confocal imaging of Raji cells using CBFB Rabbit mAb (A28470, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of RAW 264.7 cells using CBFB Rabbit mAb (A28470, dilution 1: 200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



Flow cytometry: 1X10^6 K-562 cells were intracellularly-stained with Rabbit IgG isotype control (AC042,2 µg/mL,left) or CBFB Rabbit mAb (A28470,2 µg/mL,right), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.