Leader in Biomolecular Solutions for Life Science

www.abclonal.com

NAT8 Rabbit pAb

Catalog No.: A7759

1 Publications

Basic Information

Observed MW
26kDa

Calculated MW
26kDa

Category
Polyclonal Antibodies

Applications
WB, ELISA

Cross Reactivity
Human, Mouse, Rat

Background

This gene, isolated using the differential display method to detect tissue-specific genes, is specifically expressed in kidney and liver. The encoded protein shows amino acid sequence similarity to N-acetyltransferases. A similar protein in Xenopus affects cell adhesion and gastrulation movements, and may be localized in the secretory pathway. A highly similar paralog is found in a cluster with this gene.

Recommended Dilutions

WB	1:500 - 1:2000

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
9027

Swiss-Prot
Q9UHE5

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
GLA; CML1; CCNAT; Hcml1; ATase2; TSC501; TSC510; NAT8

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG

Purity
Affinity purification

Storage
Store at -20℃

Validation Data



Western blot analysis of various lysates using NAT8 Rabbit pAb (A7759) at 1:500 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.