INADL Rabbit pAb

Catalog No.: A8476

www.abclonal.com

Basic Information

Observed MW
255 kDa

Calculated MW
196 kDa

Category
Polyclonal Antibodies

Applications
WB, ELISA

Cross Reactivity
Human

Background

This gene encodes a protein with multiple PDZ domains. PDZ domains mediate protein-protein interactions, and proteins with multiple PDZ domains often organize multimeric complexes at the plasma membrane. This protein localizes to tight junctions and to the apical membrane of epithelial cells. A similar protein in Drosophila is a scaffolding protein which tethers several members of a multimeric signaling complex in photoreceptors.

Recommended Dilutions

WB	1:500 - 1:2000

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
10207

Swiss-Prot
Q8NI35

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
Cipp; INADL; hINADL; InaD-like

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG

Purity
Affinity purification

Storage
Store at -20℃

Validation Data



Western blot analysis of various lysates using INADL Rabbit pAb (A8476) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.