Leader in Biomolecular Solutions for Life Science

ABflo® 594-conjugated Goat anti-Rabbit IgG (H+L)

Catalog No.: AS039

116 Publications

www.abclonal.com

Basic Information

Observed MW

Calculated MW

Category
Secondary Antibodies

Applications
IF/ICC, FC

Cross Reactivity
Rabbit

Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Recommended Dilutions

IF/ICC	1:100 - 1:400
FC	1:100 - 1:1000

Immunogen Information

Gene ID **Swiss-Prot**

Immunogen
This information is considered to be commercially sensitive.

Synonyms

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Goat

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃

Validation Data



Confocal imaging of HeLa cells using [KD Validated] TIMM10 Rabbit mAb (A24996, dilution 1:200) followed by a further incubation with ABflo® 594-conjugated Goat Anti-Rabbit IgG (H+L) (AS039,dilution 1:400). DAPI was used for nuclear staining (Blue). Objective: 100x.



Flow cytometry: 1X10^6 K-562 cells (negative control,left) and A-431 cells (right) were surface-stained with Purified Rabbit anti-Human E-Cadherin mAb(5 μl/Test, orange line) or secondary antibody only (blue line). Non-fluorescently stained K-562 and A-431 cells were used as blank control (red line). ABflo® 594-conjugated Goat Anti-Rabbit IgG (H+L)(AS039, 1:1000) was used as a secondary antibody.