Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Caspase-6 Rabbit mAb

Catalog No.: A19552

Recombinant | 4 Publications

Basic Information

Observed MW
33 kDa

Calculated MW
33 kDa

Category
Monoclonal Antibodies

Applications
WB, IF/ICC, ELISA

Cross Reactivity
Human, Rat

Clone No.
ARC0031

Background

This gene encodes a member of the cysteine-aspartic acid protease (caspase) family of enzymes. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic acid residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein is processed by caspases 7, 8 and 10, and is thought to function as a downstream enzyme in the caspase activation cascade. Alternative splicing of this gene results in multiple transcript variants that encode different isoforms.

Recommended Dilutions

WB	1:1000 - 1:4000
IF/ICC	1:50 - 1:200

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
839

Swiss-Prot
P55212

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
MCH2; CSP-6; caspase-6; Caspase-6

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃
Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates using Caspase-6 Rabbit mAb (A19552) at 1: 1000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90 s.



Immunofluorescence analysis of C6 cells using Caspase-6 Rabbit mAb (A19552) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.