Leader in Biomolecular Solutions for Life Science

[KO Validated] TYSY Rabbit pAb

Catalog No.: A21642

www.abclonal.com

Basic Information

Observed MW
36kDa

Calculated MW
36kDa

Category
Polyclonal Antibodies

Applications
WB, IF/ICC, IHC-P, ELISA

Cross Reactivity
Human, Mouse, Rat

Background

Thymidylate synthase catalyzes the methylation of deoxyuridylate to deoxythymidylate using, 10-methylenetetrahydrofolate (methylene-THF) as a cofactor. This function maintains the dTMP (thymidine-5-prime monophosphate) pool critical for DNA replication and repair. The enzyme has been of interest as a target for cancer chemotherapeutic agents. It is considered to be the primary site of action for 5-fluorouracil, 5-fluoro-2-prime-deoxyuridine, and some folate analogs. Expression of this gene and that of a naturally occurring antisense transcript, mitochondrial enolase superfamily member 1 (GeneID:55556), vary inversely when cell-growth progresses from late-log to plateau phase. Polymorphisms in this gene may be associated with etiology of neoplasia, including breast cancer, and response to chemotherapy.

Recommended Dilutions

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:200
IHC-P	1:50 - 1:200

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
7298

Swiss-Prot
P04818

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
TS; TMS; DKCD; HST422; SY

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Immunofluorescence analysis of C6 cells using [KO Validated] TYSY Rabbit pAb (A21642) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH-3T3 cells using [KO Validated] TYSY Rabbit pAb (A21642) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using [KO Validated] TYSY Rabbit pAb (A21642) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Mouse spleen using [KO Validated] TYSY Rabbit pAb (A21642) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Western blot analysis of lysates from wild type(WT) and TYSY knockout (KO) 293T cells using [KO Validated] TYSY Rabbit pAb (A21642) at 1:1000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.