Leader in Biomolecular Solutions for Life Science

www.abclonal.com

APRT Rabbit pAb

Catalog No.: A5456

2 Publications

Basic Information

Observed MW
22kDa

Calculated MW
20kDa

Category
Polyclonal Antibodies

Applications
WB, ELISA

Cross Reactivity
Human, Mouse

Background

Adenine phosphoribosyltransferase belongs to the purine/pyrimidine phosphoribosyltransferase family. A conserved feature of this gene is the distribution of CpG dinucleotides. This enzyme catalyzes the formation of AMP and inorganic pyrophosphate from adenine and 5-phosphoribosyl-1-pyrophosphate (PRPP). It also produces adenine as a by-product of the polyamine biosynthesis pathway. A homozygous deficiency in this enzyme causes 2,8-dihydroxyadenine urolithiasis. Two transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB	1:500 - 1:2000

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
353

Swiss-Prot
P07741

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
AMP; APRTD; APRT

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG

Purity
Affinity purification

Storage
Store at -20℃
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from 293T cells using APRT Rabbit pAb (A5456) at 1: 2500 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit (RM00021).
Exposure time: 30 s.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.