Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Occludin Rabbit mAb

Catalog No.: A24601

Recombinant | 21 Publications

Basic Information

Observed MW
16-65 kDa

Calculated MW
59kDa

Category
Monoclonal Antibodies

Applications
WB, IF-P, IHC-P, ELISA

Cross Reactivity
Human, Mouse, Rat

Clone No.
ARC63373

Background

This gene encodes an integral membrane protein that is required for cytokine-induced regulation of the tight junction paracellular permeability barrier. Mutations in this gene are thought to be a cause of band-like calcification with simplified gyration and polymicrogyria (BLC-PMG), an autosomal recessive neurologic disorder that is also known as pseudo-TORCH syndrome. Alternative splicing results in multiple transcript variants. A related pseudogene is present 1.5 Mb downstream on the q arm of chromosome 5.

Recommended Dilutions

WB	1:2000 - 1:8000
IF-P	1:200 - 1:400
IHC-P	1:200 - 1:800
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
100506658

Swiss-Prot
Q16625

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
BLCPMG; PTORCH1; PPP1R115; Occludin

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Western blot analysis of various lysates using Occludin Rabbit mAb (A24601) at 1: 2000 dilution incubated overnight at 4℃.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 20 s.




Confocal imaging of paraffin-embedded Human colon tissue using Occludin Rabbit mAb (A24601, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Human breast cancer tissue using Occludin Rabbit mAb (A24601, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Mouse colon tissue using Occludin Rabbit mAb (A24601, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using Occludin Rabbit mAb (A24601) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human liver cancer tissue using Occludin Rabbit mAb (A24601) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using Occludin Rabbit mAb (A24601) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using Occludin Rabbit mAb (A24601) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using Occludin Rabbit mAb (A24601) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.